

---

Show all appropriate work.

---

1. Write how long it took you to complete the assignment.
2. Show that for two vectors  $\mathbf{u}$  and  $\mathbf{v}$  making an angle  $\theta$  with one another

$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \cdot \|\mathbf{v}\| \cdot \cos(\theta).$$

3. Problems from the book: (I have taken these problems from the 4th edition. I have attached a scan of the book problems to this pdf.)
  - (a) Section 1.1: 1, 3, 9, 10, 12, 13, 16.
  - (b) Section 1.2: 4, 5, 12, 13, 16, 19.

## Problem Set 1.1

Problems 1–9 are about addition of vectors and linear combinations.

- 1 Describe geometrically (line, plane, or all of  $\mathbb{R}^3$ ) all linear combinations of

$$* \text{ (a) } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \text{ and } \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix} \quad \text{(b) } \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix} \quad \text{(c) } \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$$

- 2 Draw  $v = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$  and  $w = \begin{bmatrix} -2 \\ 2 \end{bmatrix}$  and  $v + w$  and  $v - w$  in a single  $xy$  plane.

- 3 If  $v + w = \begin{bmatrix} 5 \\ 1 \end{bmatrix}$  and  $v - w = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$ , compute and draw  $v$  and  $w$ .

- 4 From  $v = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$  and  $w = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ , find the components of  $3v + w$  and  $cv + dw$ .

- 5 Compute  $u + v + w$  and  $2u + 2v + w$ . How do you know  $u, v, w$  lie in a plane?

$$\text{In a plane} \quad u = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad v = \begin{bmatrix} -3 \\ 1 \\ -2 \end{bmatrix}, \quad w = \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix}.$$

- 6 Every combination of  $v = (1, -2, 1)$  and  $w = (0, 1, -1)$  has components that add to \_\_\_\_\_. Find  $c$  and  $d$  so that  $cv + dw = (3, 3, -6)$ .

- 7 In the  $xy$  plane mark all nine of these linear combinations:

$$c \begin{bmatrix} 2 \\ 1 \end{bmatrix} + d \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad \text{with } c = 0, 1, 2 \quad \text{and } d = 0, 1, 2.$$

- 8 The parallelogram in Figure 1.1 has diagonal  $v + w$ . What is its other diagonal? What is the sum of the two diagonals? Draw that vector sum.


- 9 If three corners of a parallelogram are  $(1, 1)$ ,  $(4, 2)$ , and  $(1, 3)$ , what are all three of the possible fourth corners? Draw two of them.

Problems 10–14 are about special vectors on cubes and clocks in Figure 1.4.

- 10 Which point of the cube is  $i + j$ ? Which point is the vector sum of  $i = (1, 0, 0)$  and  $j = (0, 1, 0)$  and  $k = (0, 0, 1)$ ? Describe all points  $(x, y, z)$  in the cube.

- 11 Four corners of the cube are  $(0, 0, 0)$ ,  $(1, 0, 0)$ ,  $(0, 1, 0)$ ,  $(0, 0, 1)$ . What are the other four corners? Find the coordinates of the center point of the cube. The center points of the six faces are \_\_\_\_\_.

- 12 How many corners does a cube have in 4 dimensions? How many 3D faces? How many edges? A typical corner is  $(0, 0, 1, 0)$ . A typical edge goes to  $(0, 1, 0, 0)$ .

Figure 1.4: Unit cube from  $i, j, k$  and twelve clock vectors.

- 13 (a) What is the sum  $V$  of the twelve vectors that go from the center of a clock to the hours 1:00, 2:00, ..., 12:00?
- (b) If the 2:00 vector is removed, why do the 11 remaining vectors add to 8:00?
- (c) What are the components of that 2:00 vector  $v = (\cos \theta, \sin \theta)$ ?
- 14 Suppose the twelve vectors start from 6:00 at the bottom instead of (0, 0) at the center. The vector to 12:00 is doubled to (0, 2). Add the new twelve vectors.

Problems 15–19 go further with linear combinations of  $v$  and  $w$  (Figure 1.5a).

- 15 Figure 1.5a shows  $\frac{1}{2}v + \frac{1}{2}w$ . Mark the points  $\frac{3}{4}v + \frac{1}{4}w$  and  $\frac{1}{4}v + \frac{1}{4}w$  and  $v + w$ .
- 16 Mark the point  $-v + 2w$  and any other combination  $cv + dw$  with  $c + d = 1$ . Draw the line of all combinations that have  $c + d = 1$ .
- 17 Locate  $\frac{1}{3}v + \frac{1}{3}w$  and  $\frac{2}{3}v + \frac{2}{3}w$ . The combinations  $cv + cw$  fill out what line?
- 18 Restricted by  $0 \leq c \leq 1$  and  $0 \leq d \leq 1$ , shade in all combinations  $cv + dw$ .
- 19 Restricted only by  $c \geq 0$  and  $d \geq 0$  draw the “cone” of all combinations  $cv + dw$ .




Figure 1.5: Problems 15–19 in a plane

Problems 20–25 in 3-dimensional space

## Problem Set 1.2

- 1 Calculate the dot products  $u \cdot v$  and  $u \cdot w$  and  $u \cdot (v + w)$  and  $w \cdot v$ :

$$u = \begin{bmatrix} -.6 \\ .8 \end{bmatrix} \quad v = \begin{bmatrix} 3 \\ 4 \end{bmatrix} \quad w = \begin{bmatrix} 8 \\ 6 \end{bmatrix}.$$

- 2 Compute the lengths  $\|u\|$  and  $\|v\|$  and  $\|w\|$  of those vectors. Check the Schwarz inequalities  $|u \cdot v| \leq \|u\| \|v\|$  and  $|v \cdot w| \leq \|v\| \|w\|$ .

- 3 Find unit vectors in the directions of  $v$  and  $w$  in Problem 1, and the cosine of the angle  $\theta$ . Choose vectors  $a, b, c$  that make  $0^\circ, 90^\circ$ , and  $180^\circ$  angles with  $w$ .

- 4 For any unit vectors  $v$  and  $w$ , find the dot products (actual numbers) of

(a)  $v$  and  $-v$       (b)  $v + w$  and  $v - w$       (c)  $v - 2w$  and  $v + 2w$

- 5 Find unit vectors  $u_1$  and  $u_2$  in the directions of  $v = (3, 1)$  and  $w = (2, 1, 2)$ . Find unit vectors  $U_1$  and  $U_2$  that are perpendicular to  $u_1$  and  $u_2$ .

- 6 (a) Describe every vector  $w = (w_1, w_2)$  that is perpendicular to  $v = (2, -1)$ .  
 (b) The vectors that are perpendicular to  $V = (1, 1, 1)$  lie on a \_\_\_\_\_.  
 (c) The vectors that are perpendicular to  $(1, 1, 1)$  and  $(1, 2, 3)$  lie on a \_\_\_\_\_.

- 7 Find the angle  $\theta$  (from its cosine) between these pairs of vectors:

(a)  $v = \begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix}$  and  $w = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$       (b)  $v = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix}$  and  $w = \begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix}$   
 (c)  $v = \begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix}$  and  $w = \begin{bmatrix} -1 \\ \sqrt{3} \end{bmatrix}$       (d)  $v = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$  and  $w = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$ .

- 8 True or false (give a reason if true or a counterexample if false):

- (a) If  $u$  is perpendicular (in three dimensions) to  $v$  and  $w$ , those vectors  $v$  and  $w$  are parallel.  
 (b) If  $u$  is perpendicular to  $v$  and  $w$ , then  $u$  is perpendicular to  $v + 2w$ .  
 (c) If  $u$  and  $v$  are perpendicular unit vectors then  $\|u - v\| = \sqrt{2}$ .

- 9 The slopes of the arrows from  $(0, 0)$  to  $(v_1, v_2)$  and  $(w_1, w_2)$  are  $v_2/v_1$  and  $w_2/w_1$ . Suppose the product  $v_2 w_2 / v_1 w_1$  of those slopes is  $-1$ . Show that  $v \cdot w = 0$  and the vectors are perpendicular.

- 10 Draw arrows from  $(0, 0)$  to the points  $v = (1, 2)$  and  $w = (-2, 1)$ . Multiply their slopes. That answer is a signal that  $v \cdot w = 0$  and the arrows are \_\_\_\_\_.

- 11 If  $v \cdot w$  is negative, what does this say about the angle between  $v$  and  $w$ ? Draw a 3-dimensional vector  $v$  (an arrow), and show where to find all  $w$ 's with  $v \cdot w < 0$ .

- 12 With  $v = (1, 1)$  and  $w = (1, 5)$  choose a number  $c$  so that  $w - cv$  is perpendicular to  $v$ . Then find the formula that gives this number  $c$  for any nonzero  $v$  and  $w$ . (Note:  $cv$  is the "projection" of  $w$  onto  $v$ .)
- 13 Find two vectors  $v$  and  $w$  that are perpendicular to  $(1, 0, 1)$  and to each other.
- 14 Find nonzero vectors  $u, v, w$  that are perpendicular to  $(1, 1, 1)$  and to each other.
- 15 The geometric mean of  $x = 2$  and  $y = 8$  is  $\sqrt{xy} = 4$ . The arithmetic mean is larger:  $\frac{1}{2}(x + y) = \underline{\hspace{2cm}}$ . This would come in Example 6 from the Schwarz inequality for  $v = (\sqrt{2}, \sqrt{8})$  and  $w = (\sqrt{8}, \sqrt{2})$ . Find  $\cos \theta$  for this  $v$  and  $w$ .
- 16 How long is the vector  $v = (1, 1, \dots, 1)$  in 9 dimensions? Find a unit vector  $u$  in the same direction as  $v$  and a unit vector  $w$  that is perpendicular to  $v$ .
- 17 What are the cosines of the angles  $\alpha, \beta, \theta$  between the vector  $(1, 0, -1)$  and the unit vectors  $i, j, k$  along the axes? Check the formula  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \theta = 1$ .

Problems 18–31 lead to the main facts about lengths and angles in triangles.

- 18 The parallelogram with sides  $v = (4, 2)$  and  $w = (-1, 2)$  is a rectangle. Check the Pythagoras formula  $a^2 + b^2 = c^2$  which is for *right triangles only*:

$$(\text{length of } v)^2 + (\text{length of } w)^2 = (\text{length of } v + w)^2.$$

- 19 (Rules for dot products) These equations are simple but useful:  
 (1)  $v \cdot w = w \cdot v$  (2)  $u \cdot (v + w) = u \cdot v + u \cdot w$  (3)  $(cv) \cdot w = c(v \cdot w)$   
 Use (2) with  $u = v + w$  to prove  $\|v + w\|^2 = v \cdot v + 2v \cdot w + w \cdot w$ .
- 20 The "Law of Cosines" comes from  $(v - w) \cdot (v - w) = v \cdot v - 2v \cdot w + w \cdot w$ :

$$\text{Cosine Law} \quad \|v - w\|^2 = \|v\|^2 - 2\|v\| \|w\| \cos \theta + \|w\|^2.$$


If  $\theta < 90^\circ$  show that  $\|v\|^2 + \|w\|^2$  is larger than  $\|v - w\|^2$  (the third side).

- 21 The *triangle inequality* says:  $(\text{length of } v + w) \leq (\text{length of } v) + (\text{length of } w)$ .  
 Problem 19 found  $\|v + w\|^2 = \|v\|^2 + 2v \cdot w + \|w\|^2$ . Use the Schwarz inequality  $v \cdot w \leq \|v\| \|w\|$  to show that **side 3** can not exceed **side 1** + **side 2**:

$$\text{Triangle inequality} \quad \|v + w\|^2 \leq (\|v\| + \|w\|)^2 \quad \text{or} \quad \|v + w\| \leq \|v\| + \|w\|.$$

- 22 The Schwarz inequality  $|v \cdot w| \leq \|v\| \|w\|$  by algebra instead of trigonometry:

- (a) Multiply out both sides of  $(v_1 w_1 + v_2 w_2)^2 \leq (v_1^2 + v_2^2)(w_1^2 + w_2^2)$ .  
 (b) Show that the difference between those two sides equals  $(v_1 w_2 - v_2 w_1)^2$ .  
 This cannot be negative since it is a square—so the inequality is true.



- 23 The figure shows that  $\cos \alpha = v_1/\|v\|$  and  $\sin \alpha = v_2/\|v\|$ . Similarly  $\cos \beta$  is \_\_\_\_\_ and  $\sin \beta$  is \_\_\_\_\_. The angle  $\theta$  is  $\beta - \alpha$ . Substitute into the trigonometry formula  $\cos \beta \cos \alpha + \sin \beta \sin \alpha$  for  $\cos(\beta - \alpha)$  to find  $\cos \theta = v \cdot w / \|v\| \|w\|$ .

- 24 One-line proof of the Schwarz inequality  $|u \cdot U| \leq 1$  for unit vectors:

$$|u \cdot U| \leq |u_1| |U_1| + |u_2| |U_2| \leq \frac{u_1^2 + U_1^2}{2} + \frac{u_2^2 + U_2^2}{2} = \frac{1 + 1}{2} = 1.$$

Put  $(u_1, u_2) = (.6, .8)$  and  $(U_1, U_2) = (.8, .6)$  in that whole line and find  $\cos \theta$ .

- 25 Why is  $|\cos \theta|$  never greater than 1 in the first place?
- 26 If  $v = (1, 2)$  draw all vectors  $w = (x, y)$  in the  $xy$  plane with  $v \cdot w = x + 2y = 5$ . Which is the shortest  $w$ ?
- 27 (Recommended) If  $\|v\| = 5$  and  $\|w\| = 3$ , what are the smallest and largest values of  $\|v - w\|$ ? What are the smallest and largest values of  $v \cdot w$ ?

### Challenge Problems

- 28 Can three vectors in the  $xy$  plane have  $u \cdot v < 0$  and  $v \cdot w < 0$  and  $u \cdot w < 0$ ? I don't know how many vectors in  $xyz$  space can have all negative dot products. (Four of those vectors in the plane would certainly be impossible...).
- 29 Pick any numbers that add to  $x + y + z = 0$ . Find the angle between your vector  $v = (x, y, z)$  and the vector  $w = (z, x, y)$ . Challenge question: Explain why  $v \cdot w / \|v\| \|w\|$  is always  $-\frac{1}{2}$ .
- 30 How could you prove  $\sqrt[3]{xyz} \leq \frac{1}{3}(x + y + z)$  (geometric mean  $\leq$  arithmetic mean)?
- 31 Find four perpendicular unit vectors with all components equal to  $\frac{1}{2}$  or  $-\frac{1}{2}$ .
- 32 Using  $v = \text{randn}(3, 1)$  in MATLAB, create a random unit vector  $u = v/\|v\|$ . Using  $V = \text{randn}(3, 30)$  create 30 more random unit vectors  $U_j$ . What is the average size of the dot products  $|u \cdot U_j|$ ? In calculus, the average  $\int_0^\pi |\cos \theta| d\theta / \pi = 2/\pi$ .