

Homework 5

Due Friday, October 12th at 5pm

Start reading Chapter 8 of Taylor's Classical Mechanics.

1. Taylor 7.11, p282
2. Taylor 7.14, p283
3. Taylor 7.24 & 7.27, p285
4. Consider our three-rod model for a "cat".
 - (a) Begin with all three rods aligned along the α -axis. Calculate the change in orientation, $\Delta\theta$, of the whole system along the curve in shape space that runs from $\alpha = 0$ to $\alpha = 2\pi$ along the α -axis. Draw the cat before and after the traversal of the path. [Note that because shape space is a torus, this is a closed curve. What does this mean about the shape of the cat at the end of the path?]
 - (b) Calculate $\Delta\theta$ for the same setup as in part (a) but use a path that runs along the β -axis from $\beta = 0$ to $\beta = 2\pi$. Draw the cat before and after the traversal of the path.
 - (c) Calculate the "field strength,"

$$B = \frac{\partial A_\beta}{\partial \alpha} - \frac{\partial A_\alpha}{\partial \beta}.$$

- (d) Now return to the four-legged path that we considered in class (a square of side length $\pi/2$ in shape space). Integrate the field strength B over this whole square:

$$\int_0^{\pi/2} \int_0^{\pi/2} B d\alpha d\beta.$$

Compare this result to the sum of all four $\Delta\theta$ that we calculated in class. If you have taken vector calculus explain why these two calculations compare the way that they do. If you haven't, simply write: "I haven't taken vector calculus" to get full credit.

- (e) Now that we are considering a more complicated three-rod model, the transformation to body axes can also be more complicated. For example, if we place the x'_b axis parallel to the third rod the transformation from θ to θ' will certainly depend on both α and β . In general, we can write this as

$$\theta' = \theta + \psi(\alpha, \beta),$$

where ψ is a function of both shape coordinates. Recall that we defined A_α and A_β as the coefficients of $\dot{\alpha}$ and $\dot{\beta}$ in the expression

$$\dot{\theta} = A_\alpha \dot{\alpha} + A_\beta \dot{\beta}.$$

If we switch to the new coordinate θ' , what are the expressions for the new potentials A'_α and A'_β in terms of A_α , A_β and ψ ?

- (f) Show that $B' = B$. This shows that the field strength is gauge invariant and partially explains its value. All of these calculations work similarly in E & M and other gauge theories.